







- [3] Fescina RH, De Mucio B, Martínez G, Alemán A, Sosa C, Mainero L, et al. Monitoring fetal growth- Self instruction. 2nd ed: Pan American Health Organization; 2011.
- [4] Centers for Disease Control. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Rockville, MD: U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES; 2014. p. 171.
- [5] SRNT. Biochemical verification of tobacco use and cessation. *Nicotine and Tobacco Research*, 2002;4:149-59.
- [6] Braun JM, Daniels JL, Poole C, Olshan AF, Hornung R, Bernert JT, et al. A prospective cohort study of biomarkers of prenatal tobacco smoke exposure: The correlation between serum and meconium and their association with infant birth weight. *Environmental Health*, 2010;9:53.
- [7] Wu FY, Chiu HT, Wu HD, Lin CJ, Lai JS, Kuo HW. Comparison of urinary and plasma cotinine levels during the three trimesters of pregnancy. *Paediatric and Perinatal Epidemiology*, 2008;22:296-301.
- [8] Lindqvist R, Lendahls L, Tollbom O, Aberg H, Hakansson A. Smoking during pregnancy: Comparison of self-reports and cotinine levels in 496 women. *Acta Obstetrica et Gynecologica Scandinavica*, 2002;81:240-4.
- [9] Parna K, Rahu M, Youngman LD, Rahu K, Nygard-Kibur M, Koupil I. Self-reported and serum cotinine-validated smoking in pregnant women in Estonia. *Maternal and child health journal* 2005;9:385-92.
- [10] Post A, Gilljam H, Bremberg S, Galanti MR. Maternal smoking during pregnancy: A comparison between concurrent and retrospective self-reports. *Paediatric Perinatal Epidemiology*, 2008;22:155-61.
- [11] Dejmeek J, Solansk y I, Podrazilova K, Sram RJ. The exposure of nonsmoking and smoking mothers to environmental tobacco smoke during different gestational phases and fetal growth. *Environmental Health Perspectives*, 2002;110:601-6.
- [12] Horta BL, Victora CG, Menezes AM, Halpern R, Barros FC. Low birthweight, preterm births and intrauterine growth retardation in relation to maternal smoking. *Paediatric Perinatal Epidemiology*, 1997;11:140-51.
- [13] Sclowitz IK, Santos IS, Domingues MR, Matijasevich A, Barros AJ. Maternal smoking in successive pregnancies and recurrence of low birthweight: The 2004 Pelotas birth cohort study, Brazil. *Cadernos de Saude Publica*, 2013;29:123-30.
- [14] Kharrazi M, DeLorenze GN, Kaufman FL, Eskenazi B, Bernert JT, Jr., Graham S, et al. Environmental tobacco smoke and pregnancy outcome. *Epidemiology* 2004;15:660-70.
- [15] Windham GC, Eaton A, Hopkins B. Evidence for an association between environmental tobacco smoke exposure and birthweight: a meta-analysis and new data. *Paediatric and perinatal epidemiology* 1999;13:35-57.
- [16] Calbiotech. Cotinine ELISA. Calbiotech; 2008.
- [17] Epi Info. Epi Info Tm 7. 1. 1. 14. Atlanta, GA: CDC; 2013.
- [18] Statistical Package for the Social Sciences. SPSS 21.0 for Windows. Chicago, IL2013.
- [19] Zheng Q, Wouolbright LA, Afgan I. PRAM surveillance report Alabama 2010. Montgomery, Al: Alabama Department of Public Health; 2012.
- [20] Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Menacker F, Munson ML. Births: final data for 2002. *National vital statistics reports* 2003;52:1-113.
- [21] Aurrekoetxea JJ, Murcia M, Rebagliato M, Lopez MJ, Castilla AM, Santa-Marina L, et al. Determinants of self-reported smoking and misclassification during pregnancy, and analysis of optimal cut-off points for urinary cotinine: A cross-sectional study. *BMJ open* 2013;3.
- [22] Gilligan C, Sanson-Fisher R, Eades S, Wenitong M, Panaretto K, D'Este C. Assessing the accuracy of self-reported smoking status and impact of passive smoke exposure among pregnant Aboriginal and Torres Strait Islander women using cotinine biochemical validation. *Drug and alcohol review* 2010;29:35-40.
- [23] Shipton D, Tappin DM, Vadiveloo T, Crossley JA, Aitken DA, Chalmers J. Reliability of self reported smoking status by pregnant women for estimating smoking prevalence: a retrospective, cross sectional study. *Bmj* 2009;339:b4347.
- [24] Pickett KE, Rathouz PJ, Kasza K, Wakschlag LS, Wright R. Self-reported smoking, cotinine levels, and patterns of smoking in pregnancy. *Paediatric and Perinatal Epidemiology*, 2005;19:368-76.
- [25] Webb DA, Boyd NR, Messina D, Windsor RA. The discrepancy between self-reported smoking status and urine cotinine levels among women enrolled in prenatal care at four publicly funded clinical sites. *Journal of Public Health Management and Practice*, 2003;9:322-5.
- [26] Jhun HJ, Seo HG, Lee DH, Sung MW, Kang YD, Syn HC, et al. Self-reported smoking and urinary cotinine levels among pregnant women in Korea and factors associated with smoking during pregnancy. *Journal of Korean Medical Science*, 2010;25:752-7.
- [27] West R, Zatonski W, Przewozniak K, Jarvis MJ. Can we trust national smoking prevalence figures? Discrepancies between biochemically assessed and self-reported smoking rates in three countries. *Cancer Epidemiol Biomarkers & Prevention*, 2007;16:820-2.
- [28] EPA. EPA to study formaldehyde. *Environmental building news*. Brattleboro, VT: BuildingGreen.com; 2008.
- [29] Center for Disease Control. The health consequences of involuntary smoking: A report of General Surgeon. In: Coordinating Center for Health Promotion NCfCDPaHP, editor. Atlanta, Georgia: Office on Smoking and Health; 2006.